


available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/modo

Expression of *Siamois* and *Twin* in the blastula Chordin/Noggin signaling center is required for brain formation in *Xenopus laevis* embryos

Hideyuki Ishibashi^{a,1}, Noriko Matsumura^{a,1}, Hiroshi Hanafusa^{b,c}, Kunihiro Matsumoto^{b,c}, E.M. De Robertis^d, Hiroki Kuroda^{a,*}

^aFaculty of Education (Biology), Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

^bDepartment of Molecular Biology, Graduate School of Science, Institute for Advanced Research, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan

^cCREST, Japan Science and Technology Corporation, Chikusa-ku, Nagoya 464-8602, Japan

^dHoward Hughes Medical Institute and Department of Biological Chemistry, University of California, Los Angeles, CA 90095-1662, USA

ARTICLE INFO

Article history:

Received 7 March 2007

Received in revised form

4 October 2007

Accepted 9 October 2007

Available online 12 October 2007

Keywords:

Cell signaling

Spemann organizer

Siamois

Twin

Chordin

Noggin

BMP

Wnt

Xenopus

ABSTRACT

The blastula Chordin- and Noggin-expressing (BCNE) center located in the dorsal animal region of the *Xenopus* blastula embryo contains both prospective anterior neuroectoderm and Spemann organizer precursor cells. Here we show that, contrary to previous reports, the canonical Wnt target *homeobox* genes, Double knockdown of these genes using anti-sense morpholinos in *Xenopus laevis* blocked head formation, reduced the expression of the other BCNE center genes, upregulated *Bmp4* expression, and nullified hyperdorsalization by lithium chloride. Moreover, gain- and loss-of-function experiments showed that *Siamois* and *Twin* expression is repressed by the vegetal transcription factor *VegT*. We propose that *VegT* expression causes maternal β -Catenin signals to restrict *Siamois* and *Twin* expression to the BCNE region. A two-step inhibition of BMP signals by *Siamois* and *Twin* – first by transcriptional repression of *Bmp4* and then by activation of the expression of the BMP inhibitors Chordin and Noggin – in the BCNE center is required for head formation.

© 2007 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Head formation is of considerable interest in developmental biology (reviewed in De Robertis and Kuroda, 2004; Niehrs, 2004; Stern, 2005). In vertebrate embryos, the establishment of the anteroposterior (A-P) axis has been most fully studied

in amphibians, and embryological and molecular biological approaches have led to the identification of many molecules that control A-P patterning. Several lines of evidence indicate that head formation in *Xenopus* is controlled by a maternal nuclear β -Catenin signal located in the dorsal side of the blastula embryo. However, at later stages of development zygotic

* Corresponding author. Tel./fax: +81 54 238 4304.

E-mail address: ehkurod@ipc.shizuoka.ac.jp (H. Kuroda).

¹ These authors equally contributed to this work.

0925-4773/\$ - see front matter © 2007 Elsevier Ireland Ltd. All rights reserved.

doi:10.1016/j.mod.2007.10.005

Wnt/ β -Catenin signals have the opposite effect, ventralizing the embryo (Christian and Moon, 1993). The late Wnt/ β -Catenin effects are probably mediated by the increased duration of BMP signals caused by inhibition of phosphorylation of Smad1/5/8 by GSK3 (Fuentealba et al., 2007). The early effects, however, are mediated by the well-known β -Catenin/Tcf3 (T-Cell Factor-3, a transcriptional repressor) pathway (Schneider et al., 1996; Houston et al., 2002; De Robertis and Kuroda, 2004). In the early blastula embryo, overexpression of Wnt ligands, their downstream signaling components, or their target genes, on the ventral side of the early embryo results in axis duplication, including perfect secondary head structures.

The nuclear localization of β -Catenin on the dorsal side extends broadly from the bottom (vegetal) to the top (animal) pole of the blastula in *Xenopus* (Schneider et al., 1996). The egg cytoplasm is heterogeneous, and when zygotic gene transcription starts at the mid-blastula transition the Nieuwkoop signaling center is formed in the dorsal-vegetal region (Fig. 1A). Nieuwkoop center cells express *Xenopus* Nodal-related (*Xnr1*, 2, 4, 5, and 6), TGF β -superfamily factors that have potent mesoderm-inducing activity (Agius et al., 2000; Takahashi et al., 2000). High levels of Xnrs emanating from the Nieuwkoop center induce the Spemann organizer in overlying cells at gastrula, while the Nieuwkoop center cells themselves go on to form anterior endoderm. The β -Catenin signal induces the expression of BMP antagonists such as *Chordin* and *Noggin* in the BCNE (blastula *Chordin*- and *Noggin*-expressing) center, which has only limited overlap with the Nieuwkoop center (Fig. 1A; see also Fig. 1E in Kuroda et al.,

2004). Blastula expression of *Chordin* and *Noggin* in the cellular precursors of the brain itself causes the anterior neuroectoderm to be predisposed to neural induction by the underlying endomesoderm and explains the enigmatic phenomenon of planar neural induction in dorsal Keller explants in *Xenopus* (Ruiz i Altaba, 1993; Kuroda et al., 2004).

In addition to *Chordin* and *Noggin*, other genes are expressed in a localized fashion in the BCNE center: the homeobox genes *Siamois* and *Twin*, the BMP antagonist *Xnr3*, the winged-helix gene *FoxA4a/pintallavis/HNF3 β* , and the BMP-like gene *Admp* (Fig. 1A, Kuroda et al., 2004; Wessely et al., 2004; Reversade and De Robertis, 2005). The homeobox genes *Siamois* and *Twin* are expressed immediately after the mid-blastula transition and are direct targets of the early β -Catenin/Wnt signal with potent dorsalizing activity (Lemaire et al., 1995; Carnac et al., 1996; Laurent et al., 1997). According to their high similarity in amino acids sequence (88% in the homeodomain) and perfectly overlapping expression pattern these genes may have redundant functions during *Xenopus* embryogenesis or act synergistically, possibly as heterodimers, to regulate gene transcription (Laurent et al., 1997).

In the present study, we have analyzed the function of *Siamois* and *Twin* using morpholino antisense oligomers (MO) (Heasman, 2002). The loss-of-function phenotype for *Siamois* has been previously investigated using dominant negative forms containing an engrailed repression domain (Fan and Sokol, 1997; Kessler, 1997). However, it is important to study gene function in a true loss-of-function situation by MO knockdown rather than in a dominant-negative approach, since microinjection of dominant-negative forms could affect other regions where the desired genes being investigated are not expressed, and these constructs persist much longer time in the embryo than the wild-type products. We report here that the *Siamois* and *Twin* in combination are key regulators of all genes expressed in the BCNE center, and are required for head formation and planar neural induction. Furthermore, we show that the vegetal expression of the maternal *T-box* transcription factor *VegT* (Zhang et al., 1998) is the key component that restricts the expression region of *Siamois* and *Twin* to the dorsal-animal BCNE center, thereby generating the distinction between the BCNE and Nieuwkoop signaling centers in the animal and vegetal hemispheres.




Fig. 1 – *Siamois* and *Twin* are regulated by β -Catenin. (A) The positioning of two dorsal signaling centers at blastula. The BCNE center expresses *Chordin* (*Chd*), *Noggin* (*Nog*), *Siamois* (*Sia*), *Twin* (*Twin*), *Xenopus nodal-related 3* (*Xnr3*), *FoxA4a/Pintallavis/HNF3 β* , *Anti-dorsalizing morphogenetic protein* (*ADMP*). D, dorsal; V, ventral. (B) The effect of β -Cat-MO, demonstrating that BCNE center genes require β -Catenin. (C–F) Blocking of Xnr inductive events by CerS mRNA injection inhibits *Chordin* expression at gastrula but not at blastula (when the BCNE is active), while *Siamois* and *Twin* are independent of Xnrs. *Ornithine decarboxylase* (*ODC*) was used as loading control.

2. Results and discussion

2.1. Depletion of both *Siamois* and *Twin* causes anterior truncations

The expression of *Siamois*, *Twin*, *Chordin*, and *Noggin* requires maternal β -Catenin, as shown by microinjection of a MO that inhibits β -Catenin (β -Cat-MO) translation. β -Catenin depletion completely blocked the expression of these BCNE genes in the stage 9 blastula (Fig. 1B). It has been previously reported that *Twin* expression is blocked by a dominant-negative form of *Smad4* (Nishita et al., 2000). We now investigated whether *Siamois* and *Twin* expression is regulated by Nodal. The mesoderm-inducing activity of endogenous *Xenopus* nodals (Xnrs) can be blocked by over-

expression of CerS protein, the C-terminal portion of Cerberus (Bouwmeester et al., 1996; Piccolo et al., 1996; Ramis et al., 2007) (Fig. 1C–F). The expression of *Chordin* at gastrula stage is greatly inhibited by CerS, but the initial blastula expression is not. This indicates that the early expression of *Chordin* is regulated by β -Catenin, whereas *Chordin* expression at gastrula is driven by Xnrs (Fig. 1C) (Wessely et al., 2001). However, no differences were detected in *Siamois* and *Twin* expression in CerS injected embryos (Fig. 1D and E), indicating that *Siamois* and *Twin* are independent of Xnrs and are exclusively regulated by β -Catenin. In these RT-PCR analyses the house-keeping gene *Ornithine decarboxylase* (*ODC*) was used as an mRNA loading control (Fig. 1F). We conclude from these results that *Siamois* and *Twin* are under the control of the early β -Catenin signals, but not of Xnrs.

Loss-of-function studies were carried out with *Siamois* and *Twin* antisense morpholino oligomers (Sia-MO and Twn-MO) designed to hybridize to the translation initiation sites of these genes (Fig. 2A). There are only six base differences between these MOs, yet each MO effectively blocked only the translation of its own target gene (Fig. 2B). Thus, Sia-MO and Twn-MO injected on their own served as internal specificity controls. Sia-MO blocked the secondary axis inducing activity of ventrally injected *Siamois* mRNA (Fig. 2C and D), but Twn-MO did not (Fig. 2E), and vice-versa (data not shown).




Fig. 2 – The function of *Xenopus laevis* *Siamois* and *Twin* is blocked by the antisense morpholino oligomer reagents designed in this study. (A) The MOs for *Siamois* and *Twin* target the translation initiation site. (B) In vitro translated *Siamois* and *Twin* protein were blocked only by each specific MO. (C) Secondary axes induced by ventral injection *Siamois* mRNA (5 pg); 77.3% (17/22) of embryos have secondary axes with formation of complete heads. (D) Secondary axis-inducing activity by *Siamois* mRNA is blocked by Sia-MO (4 ng); 0% (0/25) of the embryos formed secondary axes. (E) Twn-MO (4 ng) did not block the effect of *Siamois* mRNA; 78.9% (15/19) of embryos had secondary axes with heads.




Fig. 3 – *Siamois* and *Twin* are required for head formation. (A) Twn-MO injected embryo. Small eyes were detected in 10.3% of injected embryos (3/29). (B) Sia-MO injected embryo. Small eyes were detected in 87.5% of embryos (42/48). (C) Twn-MO and Sia-MO co-injected embryos. 80% of embryos (28/35) lost head structures. (D) Co-injection of Twn-MO and *Siamois* mRNA rescued the Twn-MO phenotype to normal or mild dorsalization (big eyes in 42.9% (15/35) of the embryos). (E) Sia-MO and *Twin* mRNA coinjected embryo. Small eyes were detected in 14.8% of embryos (4/27). (F) Control uninjected sibling embryo for (A–E). (G) Sagittal histological section of the embryo shown in C. (H) Sagittal histological section of embryo shown in F. (I) Whole-mount in situ hybridization analysis of the anterior neural marker *Otx2* in stage 26 embryo co-injected with *Twin*-MO and *Siamois*-MO. (J) *Otx2* expression in control embryo at stage 26. fb, forebrain; mb, midbrain; hb, hindbrain; nt, neural tube; no, notochord; he, heart.

Single Sia-MO or Twn-MO injections were without phenotypic effect, except for slight reduction in eye size (Fig. 3A and B, compare to Fig. 3F). The Twn-MO small eye phenotype was over-rescued by a small amount (1 pg) of *Siamois* mRNA (Fig. 3D), and that of Sia-MO was rescued by 1 pg of *Twin* mRNA (Fig. 3E). Importantly, when both genes were depleted together, a severe head defect phenotype was observed with high penetrance (Fig. 3C). MOs for the BCNE center gene *Chordin* (*Chd*-MO) also have some head inhibitory activity (Oelgeschlager et al., 2003), but the effect of co-injection of Sia-MO and Twn-MO was much stronger (Fig. 3C). In histological sections, forebrain development, but not hindbrain or spinal cord, was severely impaired (Fig. 3G and H). In in situ hybridizations, the anterior brain marker *Otx2* was blocked by co-injection of Sia-MO and Twn-MO (Fig. 3G–J), while the more posterior neural markers *krox20* and *HoxB9* continued to be expressed (data not shown). Notochord and heart formation still took place in embryos depleted for *Siamois* and *Twin* (Fig. 3G and H). We conclude that *Siamois* and *Twin* are required for anterior head formation in *Xenopus* development, in particular the forebrain.

2.2. *Siamois* and *Twin* function as key regulators of the BCNE center

The BCNE center is under the control of β -Catenin. Lithium chloride (LiCl) inhibits glycogen synthase kinase-3 β (GSK-3), stabilizing β -Catenin and increasing its levels in the nucleus (Klein and Melton, 1996; Schneider et al., 1996). Embryos treated at the 32-cell stage with LiCl are dorso-anteriorized, resulting in radial head structures lacking trunk–tail structures (Kao et al., 1986) (Fig. 4A). *Siamois* mRNA overexpression in the whole embryo is sufficient to mimic the LiCl-dorsalized phenotype (Lemaire et al., 1995). To test whether *Siamois* and *Twin* mediate the LiCl phenotype, loss-of-function experiments were performed. Co-injection of *Sia*-MO and *Twn*-MO prior to LiCl treatment completely blocked the dorsalizing effects of LiCl (Fig. 4A–C). This indicates that *Siamois* and *Twin* constitute one of the main gene outputs acting downstream of the β -Catenin signal.




Fig. 4 – Dorsalization by LiCl and planar neural induction require *Siamois* and *Twin*. (A) Embryo treated with LiCl at the 32-cell stage. The average of dorso-anterior index (DAI); was 9.58 (Kao et al., 1986). (B) Embryo co-injected with *Sia*-MO and *Twn*-MO prior to LiCl treatment. The average of DAI was 5.38. (C) Control embryo for (A) and (B). In all cases 8 ng of MOs and 1 pg of mRNA were injected. (D) Phenotype of dorsal-animal injection of *Sia*-MO (total 4 ng) and *Twn*-MO (total 4 ng); 76.0% (19/25) of embryos lost head structures. (E) The phenotype of dorsal-vegetal injection of *Sia*-MO and *Twn*-MO. 7.7% (2/26) of embryos lost head structures. (F) Sibling control embryo at stage 38 for (A) and (B). (G) RT-PCR analyses of the effects of *Sia*-MO and *Twn*-MO on whole embryos at stage 9 and 11. (H) Neuroectoderm (NE) region of Keller explants used for molecular studies. (I) RT-PCR analyses of the effect of *Sia*-MO, *Twn*-MO, *Chd*-MO, and *Chordin* mRNA on the NE region shown in (H).

The entire BCNE center region is derived from dorsal-animal blastomeres at the 8-cell stage, whereas the Nieuwkoop center is derived from dorsal-vegetal blastomeres (Kuroda et al., 2004). To test whether *Siamois* and *Twin* also function as Nieuwkoop center genes, we injected both MOs into dorsal-animal or dorsal-vegetal blastomeres at the 8-cell stage and found that *Siamois* and *Twin* were required for head formation in the animal blastomeres, but not in ventral ones (Fig. 4D–F). At late blastula, stage 9, the expression levels of *Chordin* and *Noggin* were significantly reduced by dorsal-animal but not by dorsal-vegetal depletion of *Siamois* and *Twin* (Fig. 4G, lanes 1–3). At mid-gastrula, stage 11, the expression levels of *Chordin* and *Noggin* were not affected by loss-of-function of *Siamois* and *Twin* (Fig. 4G, lanes 4–6). These data are consistent with the view that at blastula *Siamois* and *Twin* function exclusively in the BCNE center, activating the expression of *Chordin* and *Noggin*. At gastrula stages, however, *Chordin* and *Noggin* are regulated by Nodal-related signals emanating from Spemann organizer mesoderm.

We reported previously that planar neural induction in Keller dorsal gastrula explants requires *Chordin* (Kuroda et al., 2004). We now asked whether the expression of *Siamois* and *Twin* in the prospective neuroectoderm at blastula is responsible for the “planar” neural induction signals. To investigate this, we examined gene expressions in the neuroectodermal (NE) region in Keller sandwiches (Fig. 4H) (Keller and Danilchik, 1988; Kuroda et al., 2004). The neuroectodermal region of control Keller sandwiches expressed the anterior neural marker *Otx2* and the neuronal marker *N-tubulin* (Fig. 4I, lane 2), whereas the expression of *Otx2* was blocked by co-injection of *Sia*-MO and *Twn*-MO (Fig. 4I, lane 3), just as in the case of *Chd*-MO (Fig. 4I, lane 5). These results show that the planar neural induction observed in Keller explants requires *Siamois* and *Twin*. Microinjection of *Chordin* mRNA partially rescued the anti-neural effect of *Sia*-MO and *Twn*-MO, and completely the inhibition of planar induction by *Chd*-MO (Fig. 4I, lanes 4 and 6). We conclude from these experiments that planar neural induction requires *Siamois* and *Twin*. As described earlier, this neural induction also has a complete requirement for the expression of their downstream target *Chordin* (Kuroda et al., 2004).

In addition to *Siamois* and *Twin*, the BCNE center expresses *Chordin*, *Noggin*, *Xnr3*, *FoxA4a/pintallavis/HNF3 β* , and *Admp* (Wessely et al., 2004; Reversade and De Robertis, 2005). To investigate the relationship between these other BCNE center genes and *Siamois* and *Twin*, we examined their transcription levels at blastula stage 9. Single MOs had very weak effects, although *Sia*-MO-injected embryos had a slightly stronger phenotype than *Twin*-morphants (Fig. 5A, lanes 1–3). However, all BCNE center genes, except for *Siamois* and *Twin*, were inhibited by co-injection of *Sia*-MO and *Twn*-MO (Fig. 5A, lanes 1 and 4). In particular, *Chordin*, *Noggin*, and *FoxA4a/Pintallavis/HNF3 β* expression became undetectable (Fig. 5A, lane 4 and Fig. 5B and C). The *Xnr3* promoter region has target binding sequences for LEF/TCF and for homeodomain transcription factors (McKendry et al., 1997). Thus, the residual expression observed in the case of *Xnr3* and *Admp* might be explained by a feed-forward loop in which these genes are directly regulated by β -Catenin, as well as by *Siamois* and *Twin*.




Fig. 5 – Siamois and Twin are key regulators of the transcription of BCNE center genes. (A) RT-PCR analyses of the effects of Sia-MO, Twn-MO, or both on whole embryos. **(B)** Chordin and Noggin expression at blastula in the animal cap and marginal zone (BCNE) of normal embryos. **(C)** Co-injection of Twn-MO and Sia-MO eliminates Chordin and Noggin expression. Embryos are shown in animal view.

These results indicate that the *Siamois* and *Twin* transcriptional activators are central players in the regulation of BCNE center gene expression. They are required for the expression of Chordin, Noggin and FoxA4a/Pintallavis/HNF3 β , but not for their own expression, which depends entirely on the β -Catenin signal.

2.3. Double inhibition of BMP by Siamois and Twin by transcriptional repression and BMP antagonists

The traditional embryological view held that neural tissue forms during gastrulation, when zygotic signals from dorsal mesoderm antagonize BMPs in the overlying dorsal ectoderm (Kuroda et al., 2004). Recent studies, however, have shown that the early β -Catenin is sufficient to cause neural tissue formation in the ectoderm in the absence of underlying mesoderm, explaining earlier observations of planar neural induction in *Xenopus* (Ruiz i Altaba, 1993). Some studies have previously shown that β -Catenin inhibits BMP transcription (Baker et al., 1999; Leung et al., 2003). On the other hand, we reported that nuclear β -Catenin causes the expression of Chordin and Noggin in the BCNE center and that these BMP antagonists are required for neuralization (Wessely et al., 2001; Kuroda et al., 2004). In an attempt to reconcile these two lines of evidence, we now investigated the function of *Siamois* and *Twin* in this process.




Fig. 6 – Two-step inhibition of BMP4 by Siamois and Twin. (A) Experimental procedure. **(B)** RT-PCR analyses for the effects of *Siamois*, *Twin*, and the BMP antagonists Chordin and Noggin on *Bmp4* and its target gene *Vent1*. The nodal-related target gene *Mix1* was used to indicate that CerS was effective in blocking mesoderm and endoderm formation in these experiments. **(C)** Diagram showing a two-step model for the regulation of BMP4 via *Siamois* and *Twin*. BMP signaling is blocked in two steps (1) transcriptional repression and (2) BMP antagonists such as Chordin and Noggin which are induced by *Siamois* and *Twin*.

Small dorsal and ventral ectodermal explants were excised from embryos injected with CerS mRNA to prevent mesoderm formation (Fig. 6A). CerS, which consists of the cystine knot of Cerberus, is an excellent reagent for this type of experiment because it blocks specifically the activity of all mesoderm-inducing Nodal-related genes (Piccolo et al., 1996; Agius et al., 2000). Double depletion of *Siamois* and *Twin* caused *Bmp4* to be expressed in the dorsal animal cap fragments (Fig. 6B, compare lanes 2 and 4), and injection of *Siamois* mRNA inhibited *Bmp4* expression in ventral animal cap fragments (Fig. 6B, lanes 5 and 7). Depletion of Chordin and Noggin increased expression of the BMP target *Vent1* dorsally (Fig. 6B, lanes 2 and 3), and Chordin mRNA inhibited *Vent1* but did not affect *Bmp4* transcription (Fig. 6B; lanes 5 and 6). At the gastrula stage, the opposing homeobox genes *Gooseoid* and *Vent1/2* have been reported to be mutually antagonistic in dorsal-ventral patterning (Sander et al., 2007). We believe that *Siamois* and *Twin* act on *Vent1* at the blastula stage, before the later gastrula regulatory event of the mutual repression by *gooseoid* and *Vent1* takes place.

These results suggest to us a model in which *Siamois* and *Twin* first downregulate endogenous *Bmp4* mRNA transcription at the blastula dorsal animal region in a BMP antagonist-independent manner (as proposed by Baker et al., 1999; Leung et al., 2003). In a second step, Chordin and Noggin produced in the BCNE center inhibit any residual BMPs. In this two-step model, depicted in Fig. 6C, BMP signals are cleared

out in the BCNE center by transcriptional repression and also by BMP antagonists, both dependent on *Siamois* and *Twin* expression.

Since *Siamois* and *Twin* are transcriptional activators (Kessler, 1997), how can they repress transcription of *Bmp4*? In zebrafish, the homeobox gene *bozozok* induced by the Wnt/ β -Catenin signal encodes a transcriptional repressor and is able to directly inhibit *Bmp2b* transcription (Leung et al., 2003). A distinct possibility is that *Xenopus* might have an as yet undiscovered *bozozok*-like transcriptional repressor homologue induced by *Siamois* and *Twin* that negatively regulates the transcription levels of *Bmp4*, explaining arrow number 1 in Fig. 6C.

2.4. *VegT* defines the border between the BCNE center and the Nieuwkoop center

What determines the molecular differences between the BCNE and Nieuwkoop centers? To address this, we used a *VegT*-MO (Heasman et al., 2001) against the maternally expressed T-box transcription factor *VegT* (Zhang et al., 1998). This morpholino does not inhibit translation of the zygotic alternatively spliced *VegT* transcript designated as *Antipodean* (Stennard et al., 1999). We noted that *VegT*-MO-injected embryos always had big heads and cement glands, which are typical signs of low-BMP (Fig. 7A and B). This is what one might expect from an expansion of the BCNE center.

To investigate whether *VegT* controls BCNE center gene expression, we injected both *Wnt8* mRNA to activate β -Catenin (Christian and Moon, 1993) together with *VegT* mRNA. This combination recreates dorsal-vegetal conditions, mimicking a Nieuwkoop center in animal caps. In the conventional view currently held by the *Xenopus* field, *Siamois* and *Twin* are expressed in the Nieuwkoop center (Lemaire et al., 1995; Laurent et al., 1997). Therefore, the expectation would be that animal caps co-injected with *Wnt8* and *VegT* mRNAs would express large amounts of *Siamois* and *Twin*. Surprisingly, BCNE center genes such as *Chordin*, *Noggin*, *Siamois*, and *Twin*, which are strongly induced by increasing doses of *Wnt8* mRNA (Fig. 7C, lanes 2–5), were not upregulated in the presence of *VegT* mRNA (Fig. 7C, lanes 6–8). In contrast, the Nieuwkoop center genes *Xnr2* and *Xnr6* were induced by *VegT* mRNA (Fig. 7C, lanes 6–9). These data show that *VegT* mRNA inhibits the induction of BCNE center genes caused by *Wnt8*/ β -Catenin (Fig. 7C).

Loss-of-function experiments support the view that *VegT* serves to exclude the BCNE from the vegetal pole. The *Chordin* and *Siamois* expressing region at the BCNE center was greatly expanded towards the vegetal side by depletion of *VegT* (Fig. 7D–G). As shown earlier, this early expression of *Chordin* requires *Siamois* and *Twin* (Fig. 5A–C). This result is somewhat conflicting with a previous report by Houston et al. (2002) in which the inhibition of *VegT* reduced the level of *Chordin* and




Fig. 7 – The BCNE center is restricted to the dorsal-animal region by the maternally localized *VegT* transcription factor. (A) The phenotype of *VegT*-MO. White arrowhead indicates enlarged cement gland. (B) Control embryo for (A). (C) RT-PCR analyses of the effects of *Wnt8* and *VegT* mRNA for animal caps. 1, 4, or 16 pg of *Wnt8* mRNA and 100 pg of *VegT* mRNA were injected. (D) *Chordin* expression in control embryo at stage 9. White and black arrowheads indicate the animal and vegetal poles, respectively. (E) *Siamois* expression in control embryo. (F) *Chordin* expression of *VegT*-MO (16 ng) injected embryo. (G) *Siamois* expression of *VegT*-MO injected embryo. Note the expansion of *Chordin* and *Siamois* expression towards the vegetal side upon inhibition of *VegT*. (H) Real-time PCR analyses of *Chordin* mRNA expression levels from stage 8 to 12. (I) *Siamois* mRNA expression levels in the same samples. (J) Model of the regulation of the two blastula signaling centers. The border between the BCNE and Nieuwkoop centers is generated by repression by two maternally provided transcription factors: B1-type Sox (animal; Zhang et al., 2004) and *VegT* (vegetal).

Gooseoid, while we now report an increase. Therefore, we examined by real-time PCR analysis the level of expression of the mRNAs for *Chordin* and *Siamois* over the time course from stages 8 to 12 (Fig. 7H and I). The expression of *Chordin* and *Siamois* in VegT-MO-injected embryos was increased to approximately twice of control levels at stage 9. However, at stage 12 *Chordin* expression was greatly reduced by VegT-MO, while *Siamois* expression was not changed at stage 12 (Fig. 7H and I). One important experimental difference is that Houston et al. (2002) used antisense DNA oligo depletion in oocytes, while we used an antisense morpholino injected into cleavage stage embryos. It has been observed previously that these two loss-of-function approaches result in similar, yet not always identical phenotypes (Heasman et al., 2001). In addition, the timing of the samples may have had slight differences (Fig. 7H and I). It should also be mentioned that previous reports have shown that *Chordin* expression is activated by β -Catenin in animal but not vegetal cells (Darras et al., 1997).

Taken together, the results presented in this section suggest that *Siamois* and *Twin* are regulated differently from Nieuwkoop center genes, since VegT represses them. Recently, maternal B1-type Sox genes expressed in the animal cap have been shown to restrict *Xnr* expression, except for *Xnr3*, to the vegetal hemisphere (Zhang et al., 2004). Thus, it appears that the border between the BCNE and Nieuwkoop centers is established by the maternally encoded gene products VegT and B1-Sox. Fig. 7J shows a model in which VegT and B1-Sox are proposed to regulate the distinction between the BCNE and Nieuwkoop centers within the dorsal region of stabilized nuclear β -Catenin.

2.5. The making of a brain

The role of the prospective neuroectoderm in amphibian neural induction has been the subject of much debate (Spemann, 1938; Holtfreter et al., 1955; Nieuwkoop and Koster, 1995; Kuroda et al., 2004). It has long been known that dorsal animal cap explants respond much better than ventral ones to induction by dorsal organizer mesoderm and the mesoderm-inducer Activin (Sharpe et al., 1987; Sokol and Melton, 1991). In addition, dorsal-animal blastomeres at the 8-cell stage that have not received mesoderm induction signals, have a self-differentiation predisposition to become anterior neural tissue (Kinoshita et al., 1993).

The BCNE center is formed in cells that give rise to neuroectoderm, in particular brain, even in the absence of a mesodermal substratum. Extirpation and transplantation experiments have shown that the BCNE center is required for brain formation in the embryo (Kuroda et al., 2004). Recently, the requirement of *Noggin* expressed in the blastula animal cap ectoderm for neural differentiation has provided independent support for our proposal that the BCNE in prospective neuroectoderm plays a critical role in the differentiation of the central nervous system in *Xenopus* (Huang et al., 2007). The results of the present study show that all known BCNE center genes require the combined action of the homeobox genes *Siamois* and *Twin*. Thus, the process of brain induction can be traced back to the earliest stages of embryonic development, as is the case in other embryos such as the chick (reviewed in Stern, 2006; De Robertis and Kuroda, 2004). Neural

induction is not an event that occurs at gastrula as believed classically (Spemann, 1938), but rather starts with the cortical rotation of the 1-cell zygote, which stabilizes nuclear β -Catenin in the dorsal side of the embryo. The prospective neuroectoderm is activated by β -Catenin and repressed by VegT on the vegetal side (Fig. 7J). β -Catenin activates expression of *Siamois* and *Twin* at mid-blastula, causing the repression of *Bmp4* transcription (Baker et al., 1999; Leung et al., 2003), as well as the expression of the *Bmp*-antagonists *Chordin* and *Noggin* in the prospective neuroectoderm (Kuroda et al., 2004; Huang et al., 2007). Using morpholino knockdown we have demonstrated here that the *Siamois* and *Twin* homeobox proteins are required for the expression of BCNE center genes such as *Chordin*, *Noggin*, *FoxA4a/pintallavis/HNF3 β* , *Xnr3*, and *Admp*, which in turn participate in the formation of the vertebrate brain. In addition, we showed that *Siamois* and *Twin* are also required for the dorsalizing effects of LiCl treatment of early *Xenopus* embryos. In conclusion, *Siamois* and *Twin* represent the main output of β -Catenin in the future neuroectoderm at the blastula stage, and these homeobox genes play a key role as determinants of brain formation in *Xenopus*.

3. Experimental procedures

3.1. Embryo manipulations

Xenopus laevis embryos were obtained by *in vitro* fertilization (Sive et al., 2000). Animal cap assays were performed in 1 \times Steinberg's solution (58 mM NaCl, 0.67 mM KCl, 0.83 mM MgSO₄, 0.34 mM Ca(NO₃)₂, 4.6 mM Tris-HCl, pH 7.4, 100 mg/L Kanamycin) at stage 9 (7 h after fertilization at 22 °C). PCR conditions and primers, as well the protocol for whole-mount *in situ* hybridization, are described in <http://www.kuroda-lab.com>.

3.2. RNA and morpholino injections

To generate synthetic mRNAs, pRN3-*Siamois* was linearized with Sfi1 and transcribed with T3 RNA polymerase (Lemaire et al., 1995), pST64T-*Twin* with Sal1 and SP6 polymerase (Laurent et al., 1997), and pCS2-*Chd*, pCS2-*Wnt8*, and pCS2-*VegT* with *NotI* and SP6 RNA polymerase. MOs for *Siamois*, *Twin*, *Chordin*, *Noggin*, and *VegT* were ordered from Gene Tools, LLC. They consisted of the following sequences; Sia-MO: 5'-TTCAGCCTCATAGGTCATGTCTGTG-3' (this work), Twn-MO: 5'-TTCAGAGT CACAAGTCATGTCTGTG-3' (this work), β -Cat-MO: 5'-TTTCAAC CGTTTCCAAAGAACCAGG-3' (Kuroda et al., 2004), Chd-MO1: 5'-ACGTTGTGTGTGGTATAGTGAGGGT-3' and Chd-MO2: 5'-ACAG GATTTTGTGGTTGTCGCGAA-3' (Oelgeschläger et al., 2003), Nog-MO: 5'-TCACAAGGCACTGGGAATGATCCAT-3' (Kuroda et al., 2004), VegT-MO: 5'-CCCGACAGCAGTTTCTCATTCCAGC-3' (Heasman et al., 2001). MOs were resuspended in sterile water to a concentration of 1 mM each and then further diluted to give working solutions.

3.3. In vitro translation

Siamois and *Twin* proteins were *in vitro* translated using TnT T3/SP6 Coupled Transcription/Translation System (Promega)

in the presence of Sia-MO or Twn-MO and [³⁵S]-methionine, according to the manufacturer's instructions. Half of each reaction was analyzed by 8% SDS-PAGE and visualized by autoradiography.

Acknowledgements

We thank P. Lemaire, K.W.Y. Cho, R. Moon, C. Wylie, and T. Hayata for materials; V. Sander and members of the Kuroda laboratory for comments on the manuscript. This work was supported by the Research Reports of Uehara Memorial Foundation and the Sumitomo Foundation, NIH Grant HD21502-21, and the Howard Hughes Medical Institute.

REFERENCES

- Agius, E., Oelgeschlager, M., Wessely, O., Kemp, C., De Robertis, E.M., 2000. Endodermal Nodal-related signals and mesoderm induction in *Xenopus*. *Development* 127, 1173–1183.
- Baker, J.C., Beddington, R.S., Harland, R.M., 1999. Wnt signaling in *Xenopus* embryos inhibits *bmp4* expression and activates neural development. *Genes Dev.* 13, 3149–3159.
- Bouwmeester, T., Kim, S., Sasai, Y., Lu, B., De Robertis, E.M., 1996. Cerberus is a head-inducing secreted factor expressed in the anterior endoderm of Spemann's organizer. *Nature* 382, 595–601.
- Carnac, G., Kodjabachian, L., Gurdon, J.B., Lemaire, P., 1996. The homeobox gene *Siamois* is a target of the Wnt dorsalisation pathway and triggers organiser activity in the absence of mesoderm. *Development* 122, 3055–3065.
- Christian, J.L., Moon, R.T., 1993. Interactions between Xwnt-8 and Spemann organizer signaling pathways generate dorsoventral pattern in the embryonic mesoderm of *Xenopus*. *Genes Dev.* 7, 13–28.
- Darras, S., Marikawa, Y., Elinson, R.P., Lemaire, P., 1997. Animal and vegetal pole cells of early *Xenopus* embryos respond differently to maternal dorsal determinants: implications for the patterning of the organiser. *Development* 124, 4275–4286.
- De Robertis, E.M., Kuroda, H., 2004. Dorsal-ventral patterning and neural induction in *Xenopus* embryos. *Annu. Rev. Cell Dev. Biol.* 20, 285–308.
- Fan, M.J., Sokol, S.Y., 1997. A role for *Siamois* in Spemann organizer formation. *Development* 124, 2581–2589.
- Fuentealba, L.C., Eivers, E., Ikeda, A., Hurtado, C., Kuroda, H., Pera, E.M., De Robertis, E.M., in press. Integrating patterning signals: Wnt/GSK3 regulates the duration of the BMP/Smad1 signal. *Cell*. doi:10.1016/j.cell.2007.09.027.
- Heasman, J., 2002. Morpholino oligos: making sense of antisense? *Dev. Biol.* 243, 209–214.
- Heasman, J., Wessely, O., Langland, R., Craig, E.J., Kessler, D.S., 2001. Vegetal localization of maternal mRNAs is disrupted by VegT depletion. *Dev. Biol.* 240, 377–386.
- Holtfreter, J., Hamburger, V., 1955. Amphibians. In: Willer, B.H. et al. (Eds.), *Analysis of Development*. Haffer, NY, pp. 230–296.
- Houston, D.W., Kofron, M., Resnik, E., Langland, R., Destree, O., Wylie, C., Heasman, J., 2002. Repression of organizer genes in dorsal and ventral *Xenopus* cells mediated by maternal XTcf3. *Development* 129, 4015–4025.
- Huang, S., Yan, B., Sullivan, S.A., Moody, S.A., 2007. Noggin signaling from *Xenopus* animal blastomere lineages promotes a neural fate in neighboring vegetal blastomere lineages. *Dev. Dyn.* 236, 171–183.
- Kao, K.R., Masui, Y., Elinson, R.P., 1986. Lithium-induced respecification of pattern in *Xenopus laevis* embryos. *Nature* 322, 371–373.
- Keller, R., Danilchik, M., 1988. Regional expression, pattern and timing of convergence and extension during gastrulation of *Xenopus laevis*. *Development* 103, 193–209.
- Kessler, D.S., 1997. *Siamois* is required for formation of Spemann's organizer. *Proc. Natl. Acad. Sci. USA* 94, 13017–13022.
- Kinoshita, K., Bessho, T., Asashima, M., 1993. Competence prepattern in the animal hemisphere of the 8-cell-stage *Xenopus* embryo. *Dev. Biol.* 160, 276–284.
- Klein, P.S., Melton, D.A., 1996. A molecular mechanism for the effect of lithium on development. *Proc. Natl. Acad. Sci. USA* 93, 8455–8459.
- Kuroda, H., Wessely, O., De Robertis, E.M., 2004. Neural induction in *Xenopus*: requirement for ectodermal and endomesodermal signals via Chordin, Noggin, β -Catenin, and Cerberus. *PLoS Biol.* 2, 623–634.
- Laurent, M.N., Blitz, I.L., Hashimoto, C., Rothbacher, U., Cho, K.W., 1997. The *Xenopus* homeobox gene *twin* mediates Wnt induction of *goosecoid* in establishment of Spemann's organizer. *Development* 124, 4905–4916.
- Lemaire, P., Garrett, N., Gurdon, J.B., 1995. Expression cloning of *Siamois*, a *Xenopus* homeobox gene expressed in dorsal-vegetal cells of blastulae and able to induce a complete secondary axis. *Cell* 81, 85–94.
- Leung, T., Bischof, J., Soll, I., Niessing, D., Zhang, D., Ma, J., Jackle, H., Driever, W., 2003. *bozozok* directly represses *bmp2b* transcription and mediates the earliest dorsoventral asymmetry of *bmp2b* expression in zebrafish. *Development* 130, 3639–3649.
- McKendry, R., Hsu, S.C., Harland, R.M., Grosschedl, R., 1997. LEF-1/TCF proteins mediate wnt-inducible transcription from the *Xenopus* nodal-related 3 promoter. *Dev. Biol.* 192, 420–431.
- Niehrs, C., 2004. Regionally specific induction by the Spemann–Mangold organizer. *Nat. Rev. Genet.* 5, 425–434.
- Nieuwkoop, P.D., Koster, K., 1995. Vertical versus planar induction in amphibian early development. *Dev. Growth Differ.* 37, 653–668.
- Nishita, M., Hashimoto, M.K., Ogata, S., Laurent, M.N., Ueno, N., Shibuya, H., Cho, K.W., 2000. Interaction between Wnt and TGF-beta signalling pathways during formation of Spemann's organizer. *Nature* 403, 781–785.
- Oelgeschlager, M., Kuroda, H., Reversade, B., De Robertis, E.M., 2003. Chordin is required for the Spemann organizer transplantation phenomenon in *Xenopus* embryos. *Dev. Cell* 4, 219–230.
- Piccolo, S., Agius, E., Leyns, L., Bhattacharyya, S., Grunz, H., Bouwmeester, T., De Robertis, E.M., 1996. The head inducer Cerberus is a multifunctional antagonist of Nodal, BMP and Wnt signals. *Nature* 397, 707–710.
- Ramis, J.M., Collart, C., Smith, J.C., 2007. Xnrs and activin regulate distinct genes during *Xenopus* development: activin regulates cell division. *PLoS ONE* 2, e213.
- Reversade, B., De Robertis, E.M., 2005. Regulation of ADMP and BMP2/4/7 at opposite embryonic poles generates a self-regulating morphogenetic field. *Cell* 123, 1147–1160.
- Ruiz i Altaba, A., 1993. Induction and axial patterning of the neural plate: planar and vertical signals. *J. Neurobiol.* 17, 233–243.
- Sander, V., Reversade, B., De Robertis, E.M., 2007. The opposing homeobox genes *Goosecoid* and *Vent1/2* self-regulate *Xenopus* patterning. *EMBO J.* 26, 2955–2965.
- Schneider, S., Steinbeisser, H., Warga, R.M., Hausen, P., 1996. β -Catenin translocation into nuclei demarcates the dorsalizing centers in frog and fish embryos. *Mech. Dev.* 57, 191–198.

- Sharpe, C.R., Fritz, A., De Robertis, E.M., Gurdon, J.B., 1987. A homeobox-containing marker of posterior neural differentiation shows the importance of predetermination in neural induction. *Cell* 50, 749–758.
- Sive, H.L., Grainger, R.M., Harland, R.M., 2000. *Early Development of Xenopus laevis*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sokol, S., Melton, D.A., 1991. Pre-existent pattern in *Xenopus* animal pole cells revealed by induction with activin. *Nature* 351, 409–411.
- Spemann, H., 1938. *Embryonic Development and Induction*. Yale University Press, New Haven, CT.
- Stennard, F., Zorn, A.M., Ryan, K., Garrett, N., Gurdon, J.B., 1999. Differential expression of VegT and Antipodean protein isoforms in *Xenopus*. *Mech. Dev.* 86, 87–98.
- Stern, C.D., 2005. Neural induction: old problem, new findings, yet more questions. *Development* 132, 2007–2021.
- Stern, C.D., 2006. Neural induction: 10 years on since the ‘default model’. *Curr. Opin. Cell Biol.* 18, 692–697.
- Takahashi, S., Yokota, C., Takano, K., Tanegashima, K., Onuma, Y., Goto, J., Asashima, M., 2000. Two novel nodal-related genes initiate early inductive events in *Xenopus* Nieuwkoop center. *Development* 127, 5319–5329.
- Wessely, O., Agius, E., Oelgeschläger, M., Pera, E.M., De Robertis, E.M., 2001. Neural induction in the absence of mesoderm: β -catenin-dependent expression of secreted BMP antagonists at the blastula stage in *Xenopus*. *Dev. Biol.* 234, 161–173.
- Wessely, O., Kim, J.I., Geissert, D., Tran, U., De Robertis, E.M., 2004. Analysis of Spemann organizer formation in *Xenopus* embryos by cDNA macroarrays. *Dev. Biol.* 269, 552–566.
- Zhang, J., Houston, D.W., King, M.L., Payne, C., Wylie, C., Heasman, J., 1998. The role of maternal VegT in establishing the primary germ layers in *Xenopus* embryos. *Cell* 94, 419–421.
- Zhang, C., Basta, T., Hernandez-Lagunas, L., Simpson, P., Stemple, D.L., Artinger, K.B., Klymkowsky, M.W., 2004. Repression of nodal expression by maternal B1-type SOXs regulates germ layer formation in *Xenopus* and zebrafish. *Dev. Biol.* 273, 23–37.